Pfeiffer RF, 63
Pinner G, 142
Pollak CP, 168

Rafalson L, 114
Ranieri P, 75
Reisberg B, 22, 93
Relkin N, 114
Rogaeva EA, 1
Rordorf G, 51
Rozzini R, 75

Sage S, 58
St. George-Hyslop PH, 1
Salloway S, 79
Schmahmann JD, 51
Schwarzschild M, 51
Sharrow NA, 161
Sheikh J, 58
Shores M, 158
Sample LC, 99
Souren LEM, 22, 93
Spain W, 158
Spigler M, 114
Steffes S, 119
Stokes PE, 168
Swartz JR, 67, 154

Tannenhaus J, 146
Tariot PN, 130
Teunisse S, 33
Tinklenberg J, 58
Torossian CL, 22
Trabucchi M, 39, 75
Tsuang D, 158
Tsuda T, 1

van Crevel H, 33
van Gool WA, 33
Voci JM, 99

Wagner DR, 168
Walstra GJM, 33
Wertheimer J, 105
Westlake R, 79
Whitehouse PJ, 99
Wicks D, 58
Widman LP, 63
Wohl M, 67
Workman RH, 7, 55
Wormstall H, 127

Yesavage JA, 58

Zarcone V, 58

Subject Index

Actigraph measures, 58–62
Activities of daily living, 146–153
Adaptive kitchen behavior, 146–153
Affective disorders, 105–109, 158–160
Agitation, 95, 96–97, 175
Alzheimer, Alois, 127–129
Alzheimer's disease, 1–5, 15–20, 22–26, 29–31, 33–38, 49–50, 58–62, 67–73, 93–97, 99–103, 133, 154–156, 161–167, 175
Antidepressants, 174
Antidiuretic hormone, 75–78
Anxiety, 95

Behavioral problems, 58–62, 154–156, 168–173
Bereavement, 47
Bipolar affective disorder, 105–109, 158–160
Blessed Dementia Scale, 161, 162
Brain syndromes, 111–113
Brief Psychiatric Rating Scale, 7–10
Brief Visuospatial Memory Test, 148

Carcinoma, 111–113
CARE depression scale, 169, 170
Caregivers, 168–173
Cerebral energy metabolism, 29–31
Clinical Dementia Rating Scale, 161–167
Clozapine, 63–66, 89
Cognitive function, 161–162, 165, 166
Cohen-Mansfield Agitation Inventory, 7–10
Community elders, 168–173
Compulsive behaviors, 154–156
Continence, 22–26
Cortical dementia, 82–83
Cortical infarction, 84

Delirium, 7–10
Delusions, 88
Dementia, 15–20, 51–57, 67–73, 82–83, 114–125, 130, 154–156, 161–167
Depression, 7–10, 47, 67–73, 89–90, 105–109, 169, 170, 175
Derogatis Affects Balance Scale, 148–149
Developmental reflexes, 22–26
Disruptive nocturnal behaviors, 168–173
Dopamine, 83–84
Down syndrome, 99–103

Drugs, 63–66, 75–78, 83–84, 89, 99–103, 119–125, 131–132, 175

Energy metabolism, 29–31
Epilepsy, 158–160

Familial Alzheimer's disease, 1–5
Frontal-complex partial status epilepticus, 158–160
Frontotemporal dementia, 67–73, 154–156

Genetics, 161–167
Geriatric Depression Scale, 148
Geriatric psychiatry outpatient clinic, 136–141
Global Assessment of Functioning, 55–57
Grasping, 95–97

Hallucinations, 11–13
Hamilton Depression Rating Scale, 7–10
Hopkins Verbal Learning Test, 148
Huntington's disease, 39–45
Hydrocephalus, 15–20, 51–54
Hyponatremia, 75–78

Idiopathic musical hallucinations, 11–13
Illusion of visitors, 79–87
Incontinence, 22–26
Instrumental activities of daily living, 146–153

Kitchen behavior, 146–153
Kitchen Skills Assessment, 149–153
Klüver-Bucy syndrome, 111–113

Late-life depression, 67–73
Late-onset Huntington's disease, 39–45
Levodopa, 63–66
Lewy body disease, 133

Manic-depressive illness, 105–109, 158–160
Mattis Dementia Rating Scale, 114–118, 169, 170
Memory clinic, 33–38
Metabolism, 29–31
Microvascular disease, 84–85
Mini-Mental State Examination, 7–10, 159–160, 161, 162
Motor activity, 168–173
Musical hallucinations, 11–13

Nasopharyngeal carcinoma, 111–113
Neurodegeneration, 29–31
Neuroleptics, 119–125
Neuromotor function, 93–97
NIMH Genetics Initiative, 161–167
Nocturnal behaviors, 168–173
Normal-pressure hydrocephalus, 15–20, 51–54

Obsessive-compulsive disorder, 154–156
Outpatient clinic, 136–141

Paratonia, 94–97
Parkinson's disease, 29–31, 63–66, 89
Partial status epilepticus, 158–160
Phenotypic heterogeneity, 1–5
Pisa syndrome, 130
Pregnancy delusions, 88
Psychiatry outpatient clinic, 136–141
Psychosis, 63–66, 89, 175
Psychotropics, 131–132

Radiation brain injury, 111–113
Rapid-cycling affective disorder, 105–109
Rating Scale for Side Effects, 7–10
Reflexes, 22–26, 95
Reset osmostat syndrome, 75–78
Right hemisphere dysfunction, 84–85

Selective serotonin-reuptake inhibitors, 99–103
Sexual function, 131–132
Single-photon emission computed tomography, 15–20
Single photon emission computerized tomography, 15–20
Sleep, 29–31, 59
Slow-wave sleep, 29–31
SPECT, 15–20
Status epilepticus, 158–160
Structured Telephone Interview for Dementia Assessment, 161–167
Subcortical dementia, 83
Suicide, 89–90
Syndrome of inappropriate antidiuretic hormone secretion, 75–78

Technetium 99-HMPAO SPECT, 15–20
Telephone Interview for Cognitive Status, 161–162, 165, 166
Temporal arteritis, 142–144
Time-based Behavioral Disturbance Questionnaire, 58–62

Venlafaxine, 75–78
Visitors, illusion of, 79–87
Visual impairment, 80–82